



A service of the National Library of Medicine
and the National Institutes of Health

My NCBI
[Sign In] [Register]

All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books

Search PubMed for [] Go Clear

Limits Preview/Index History Clipboard Details

Display AbstractPlus Show 20 Sort by Send to

All: 1 Review: 0

☐ 1: Metab Eng, 2000 Oct;2(4):339-48.

ELSEVIER
FULL-TEXT ARTICLE

Links

Directed evolution of toluene dioxygenase from *Pseudomonas putida* for improved selectivity toward cis-indandiol during indene bioconversion.

Zhang N, Stewart BG, Moore JC, Greasham RL, Robinson DK, Buckland BC, Lee C.

Department of Bioprocess R&D, Merck Research Laboratories, Rahway, New Jersey 07065, USA.

Toluene dioxygenase (TDO) from *Pseudomonas putida* F1 converts indene to a mixture of cis-indandiol (racemic), 1-indenol, and 1-indanone. The desired product, cis-(1S,2R)-indandiol, is a potential key intermediate in the chemical synthesis of indinavir sulfate (Crixivan), Merck's HIV-1 protease inhibitor for the treatment of AIDS. To reduce the undesirable byproducts 1-indenol and 1-indanone formed during indene bioconversion, the recombinant TDO expressed in *Escherichia coli* was evolved by directed evolution using the error-prone polymerase chain reaction (epPCR) method. High-throughput fluorometric and spectrophotometric assays were developed for rapid screening of the mutant libraries in a 96-well format. Mutants with reduced 1-indenol by-product formation were identified, and the individual indene bioconversion product profiles of the selected mutants were confirmed by HPLC. Changes in the amino acid sequence of the mutant enzymes were identified by analyzing the nucleotide sequence of the genes. A mutant with the most desirable product profile from each library, defined as the most reduced 1-indenol concentration and with the highest cis-(1S,2R)-indandiol enantiomeric excess, was used to perform each subsequent round of mutagenesis. After three rounds of mutagenesis and screening, mutant 1C4-3G was identified to have a threefold reduction in 1-indenol formation over the wild type (20% vs 60% of total products) and a 40% increase of product (cis-indandiol) yield.

PMID: 11120645 [PubMed - indexed for MEDLINE]

Related Links

Development of a bioconversion process for production of cis-1S,2R-indandiol from indene by recombinant *Escherichia coli* construct [Appl Microbiol Biotechnol. 1999]

Microbial conversion of indene to indandiol: a key intermediate in the synthesis of CRIVAN [Metab Eng. 1999]

Indene bioconversion by a toluene inducible dioxygenase of *Rhodococcus ruber* [Appl Microbiol Biotechnol. 2004]

Desaturation, dioxygenation, and monooxygenation reactions catalyzed by naphthalene dioxygenase from *Pseudomonas* sp. strain 9816-4. [J Bacteriol. 1995]

Engineering an indene bioconversion process for the production of cis-aminoindanol: a model system for the production of indinavir sulfate [Appl Microbiol Biotechnol. 2002]

See all Related Articles...

Display AbstractPlus Show 20 Sort by Send to

Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services